

ABELIAN CHABAUTY

DAVID ZUREICK-BROWN

ABSTRACT. These are expanded lecture notes for a series of four lectures at the Arizona Winter School on “Nonabelian Chabauty”, held March 7-11, 2020 in Tucson, Arizona.

Last update: March 16, 2020

CONTENTS

1. Introduction	1
1.1. A roadmap to Chabauty	2
1.2. Course outline, and how to read these notes	3
1.3. Acknowledgements	4
2. Abelian Chabauty	4
2.2. Computational aspects: an exercise	5
3. The uniformity conjecture	6
3.5. Evidence and records	7
3.6. Chabauty–Coleman bounds	7
3.8. A new hope	8
4. Bad Reduction	8
4.2. A few examples	9
5. Rank Favorable bounds	10
5.3. Stoll’s proof of Theorem 5.1	11
5.4. The rank of a divisor	12
5.6. Chip firing and the rank of a divisor on a graph	13
5.12. Semicontinuity of specialization	15
5.14. Rank favorable bounds for curves with totally degenerate reduction	16
5.15. Refined ranks	16
6. Tropical Geometry, Berkovich spaces, and Chabauty	16
References	17

1. INTRODUCTION

Let K be a number field and let X/K be a nice¹ curve of genus $g > 1$ whose Jacobian has rank $r := \text{rank Jac}_X(K)$.

Date: March 16, 2020.

¹smooth, projective, and geometrically integral

The *Method of Chabauty–Coleman* (alternatively: “Chabauty’s method”, “Abelian Chabauty”, or just plain, vanilla, “Chabauty”) is among the most successful and widely applicable techniques for analyzing (either theoretically or explicitly) the set $X(K)$ of K -points of a low rank ($r < g$) curve X , and is an essential part of the “explicit approaches to rational points” toolbox. In particular, with some luck, Chabauty’s method allows one to

- explicitly determine, with proof, the set $X(K)$, or
- determine an upper bound on $\#X(K)$.

Mordell conjectured in the 20’s that the set $X(K)$ of K -points of X was finite. This was famously proved in the 80’s by Faltings [Fal86] (for which he was awarded a Fields medal), with subsequent independent proofs by Vojta and Bombieri [Voj91, Bom90].

Chabauty [Cha41], building on a idea of Skolem [Sko34], gave the first substantial progress toward Mordell’s conjecture. Chabauty’s method, which used p -adic techniques to produce p -adic “locally analytic” functions, relies on the hypothesis that $r < g$. This technique sat somewhat dormant until the 80’s, when Coleman’s seminal paper [Col85] resurrected and improved Chabauty’s idea.

Machine Computation. Computer aided computational tools took some time to catch up. The first big bottleneck was to improve the techniques for compute ranks of Jacobians of curves. To execute Chabauty’s method for a particular, explicit curve, one needs to know the rank r of its Jacobian (to check the $r < g$ condition), and a basis for the Mordell–Weil group of $\text{Jac}(K)$ (or at least a finite index subgroup). An early highlight is due to Gordon and Grant [GG93]; building on work of Cassels and Flynn [Fly90, Cas89, Cas83] they work out the special case of two-descent on the Jacobian of a genus 2 hyperelliptic curve with rational Weierstrass points, and (with the help of a SUN Sparcstation) provably compute the rank of a couple examples.

This ushered in a golden era of computer assisted approaches to rational points on curves (and higher dimensional varieties). Even today there are substantial conceptual and practical improvements; one recent highlight is [BPS16], which greatly expands our abilities to compute ranks of Jacobians of non-hyperelliptic curves.

Applications. It is worth highlighting a few of the numerous applications of Chabauty’s method.

- McCallum made substantial progress towards a proof of Fermat’s Last Theorem, via a careful study of certain quotients of the Fermat curve $x^p + y^p = z^p$; for instance, [McC94] proves the “second case” of Fermat’s Last Theorem for regular primes;
- arithmetic statistics: Poonen and Stoll use Chabauty’s method to prove that 100% of odd degree hyperelliptic curves have only one rational point [PS14];
- analysis of rational points on modular curves, and Mazur’s “Program B” [RZB15];
- resolution of various generalized Fermat equations [PSS07].

1.1. A roadmap to Chabuauty.

If you’re literally starting from scratch. McCallum–Poonen, Magma exercise, Coleman’s proof (and maybe original paper).

Uniform Bounds. KRZB,KZB, Stoll, LT, Stoll (earlier). Gunther–Morrow, Park, REU paper

Uniformity conjecture. Mazur’s question; original papers; surveys.

Applications of Chabauty. Applications: Numerous papers by Bruin, Stoll, Stoll et al (4 authored iwasaw theory) etc.. 2,3,7. Poonen–Stoll; Program B; Generalized Fermat; McCallum

More on explicit approaches to rational points. Surveys: Poonen, more Poonen, Stoll, others? Books? Applications: Numerous papers by Bruin, Stoll, Stoll et al (4 authored iwasaw theory) etc.. 2,3,7. Technology: Siksek, box, elliptic Chabauty. Bruin and étale descent Higher dimensional starting point.

Do I want to separate Chabauty like techniques from non-Chabauty? E.g., étale descent vs symmetric power chabauty.

Ranks and Selmer groups. Surveys: Stoll? Silverman’s book; old papers by e.g. Swinnerton–Dyer. Poonen, Stoll, Schaefer, 237 paper, developed into BPS.

p -adic integration. Papers by Coleman. Besser, Balakrishnan. Dig for others, see my own papers. KRZB, and Utah survey. Digress on

Nonabelian Chabauty: foundations. Kim’s papers. Besser, Balakrishnan, Dogra, others. Ishai, Francis Brown, Noam Kantor, David Corwin.

Nonabelian Chabauty: Applications . Cursed curve, Samir, and Dogra S; other papers by Jen et al.

Non Chabauty. Bas, Lawrence–Venkatesh, heights (Habegger)

1.2. **Course outline, and how to read these notes.** There is already an excellent survey by McCallum and Poonen [MP12]. This is short (16 pages) and a great entry point. I recommend my survey with Katz and Rabinoff [KRZB16] for the connections between p -adic and tropical techniques, and the survey [BJ15] by Baker and Jensen for a more geometric and combinatorial perspective. I also recommend attempting the computational Exercise 2.3 below (even if one is ultimately most interested in theory).

These notes will focus on the ideas from [KZB13] and [KRZB], and the papers [LT02, Sto06, Sto19, Bak08] which inspired our work. In particular, my discussion of the foundations of Abelian Chabauty, and discussion of tropical techniques, will be abridged, and these notes are somewhat of an advertisement for [MP12] and [KRZB16].

Additionally, while reading these notes, we also recommend attempting Exercise 2.3 from Subsection 2.2, which will help to quickly come up with speed with how to perform Chabauty’s method in Magma.

Abelian Chabauty. We will start with a ‘black box’ discussion of the method of Chabauty and Coleman, addressing various points of view; this section is mostly an abridged version of [MP12], and it is recommended to read their survey alongside this section and before reading future sections.

Exhibiting Abelian Chabauty as a special case of Nonabelian Chabauty is not completely straightforward. These notes do not address this, and we instead recommend Poonen’s excellent set of notes available at

http://www-math.mit.edu/~poonen/papers/p-adic_approach.pdf.

Bad reduction. One avenue to improve on Coleman’s bound is to generalize the framework of Chabauty and Coleman’s arguments to the case of bad reduction. We will discuss the advantage of working at bad primes and the difficulties and tradeoffs that arise, starting with the work of Lorenzini–Tucker [LT02].

Rank favorable bounds. When the rank is strictly smaller than $g - 1$, there are more “inputs” to Chabauty’s method and one expects this extra flexibility to lead to improvements to the method, giving rise to “rank favorable” bounds. We’ll discuss the setup, and the translation to the notion of a “rank” of a divisor (due to Stoll [Sto06]). For a curve with good reduction, this notion of rank will be the classical one, and the improved bounds will follow from Clifford’s Theorem [Har77, Theorem IV.5.4]. In the case of bad reduction, reducible reduction, ranks are no longer as well behaved; instead, we introduce Baker’s notion of “numerical rank” [Bak08] and explain how to repair Stoll’s argument in the special case of a curve with totally degenerate reduction.

Tropical Geometry and Berkovich spaces. For a curve with bad reduction at a prime p , it had been well understood that “monodromy” and “analytic continuation” of p -adic integrals was an issue. Coleman proved that in the case of good reduction, there is no “monodromy” and the various ways of analytically continuing p -adic integrals all coincide. In the case of bad reduction, they generally do not coincide (we will discuss a simple example which illustrates this).

Stoll [Sto19] discovered that, while choices of analytic continuation genuinely do differ, they do so in a fairly controlled manner (linear, even), and was able to exploit this to prove a uniformity result for *hyperelliptic* curves of small rank.

These results all argue in the framework of rigid geometry (in the sense of Tate). Great clarification arose from the systematic reformulation via Berkovich spaces, which fill in the “missing” points of rigid spaces and which, at least in the case of curves, are fairly concrete and manageable topological spaces (they’re even Hausdorff). I’ll discuss Chabauty in the setting of Berkovich and tropical geometry and explain how modern tools (e.g., Berkovich’s contraction theorem and Thuillier’s slope formula, expounded in [BPR13]) give a clean explanation of Coleman’s “good reduction” theorem, and will discuss my work with Katz and Rabinoff [KRZB] which give uniform bounds for arbitrary (but still small rank) curves.

1.3. Acknowledgements. The author would like to thank Enis Kaya, Jackson Morrow, Dino Lorenzini, and John Voight for useful comments and/or discussions. The author was supported by National Science Foundation CAREER award DMS-1555048.

2. ABELIAN CHABAUTY

We give here a quick “black box” version of Chabauty’s method, broken into 3 parts: setup, local analysis, and global coordination. We refer the reader to the excellent survey [MP12] for a more detailed introduction.

As before, let K be a number field with ring of integers \mathcal{O}_K and let X/K be a nice curve of genus $g > 1$. Let $r := \text{rank Jac}_X(K)$ be the rank of the Jacobian of X . Fix a prime p and a prime \mathfrak{p} of \mathcal{O}_K above p .

Setup. Under the assumption $r < g$, there exist **locally analytic** functions f_ω on $X(K_p)$ (arising as a **p -adic integral** of a differential ω) which vanish on $X(K)$, but not on $X(K_p)$. More precisely, there exists a subspace $V \subset H^0(X_{K_p}, \Omega_X^1)$ such that $\dim_{K_p} V \geq g - r$, and with the property that the p -adic integral

$$\int_P^Q \omega$$

vanishes for all $P, Q \in X(K)$ and for all $\omega \in V$. We will frequently refer to V as V_{chab} . This is (more or less) enough to conclude finiteness (and is roughly the original argument of Chabauty [Cha41]). See [MP12, Section 4 and Subsection 5.4] for proofs of these statements (culminating in [MP12, Theorem 4.4]).

Local analysis. On (residue) discs, the integrals f_ω are “locally analytic”: they have (p -adic) power series expansions, a discrete set of zeroes, and are amenable to fairly explicit study via tools from p -adic analysis (Newton polygons, or in more complicated situations, tropical geometry). In Coleman’s original analysis ([Col85, Lemma 3] or [MP12, Lemma 5.1]), one can bound the number zeros of f_ω in a residue disc in terms of the zeroes of its “derivative”, which we summarize as a ‘ **p -adic Rolle’s theorem**’ (in the sense of freshman calculus). In the simplest case one gets Rolle’s theorem on the nose: for $K = \mathbb{Q}$ and $p > 2$, Coleman proves [MP12, Theorem 5.3(1)] that the number of zeroes of f_ω in a residue disc D_P is at most $1 + n_P$, where

$$n_P = \#(\text{div } \omega \cap D_P). \tag{2.0.1}$$

See [MP12, Section 5] for proofs of these statements (culminating in [MP12, Theorem 5.5]).

Remark 2.1. An exciting “modern” version of this argument is [BD19, Section 4], where they compare the divisor of a locally analytic function F to the divisor of $\mathcal{D}(F)$, where \mathcal{D} is a “nice” differential operator \mathcal{D} .

Global coordination. One needs some way to coordinate the different, a priori independent, local bounds (as in Equation 2.0.1), and typically exploits some type of “global” theorem from the geometry of curves. In Coleman’s proof, Riemann–Roch [Har77, Theorem IV.1.3] suffices; the local bounds (under the $K = \mathbb{Q}$ and $p > 2$ hypotheses) are $1 + n_P$; by Equation 2.0.1 we have that

$$\sum_{P \in X(\mathbb{F}_p)} n_P = \sum_{P \in X(\mathbb{F}_p)} \#(\text{div } \omega \cap D_P) \leq \deg \text{div } \omega = 2g - 2,$$

which suffices to prove Coleman’s theorem:

$$\#X(\mathbb{Q}) \leq \sum_{P \in X(\mathbb{F}_p)} (1 + n_P) = \sum_{P \in X(\mathbb{F}_p)} 1 + \sum_{P \in X(\mathbb{F}_p)} n_P \leq X(\mathbb{F}_p) + 2g - 2.$$

In the “improvements” to this theorem that we discuss in these notes, one instead needs some other global theorem, e.g., Clifford’s theorem (or Riemann–Roch and Clifford’s theorem for graphs, or for arithmetic curves, or for other refined rank functions). In [KRZB], which uses (in a sense) the full power of the tools from tropical geometry, this step relies global information about sections of the “tropical canonical bundle” (see [KRZB, Lemma 4.15]).

Again, please see [MP12] (especially the detailed examples in Section 8) for a survey and a more thorough introduction. It is also very useful to attempt the Magma exercise (Exercise 2.3) described in the “Computational aspects” part of Subsection 1.2.

2.2. Computational aspects: an exercise. While reading these notes, we also recommend attempting Exercise 2.3 below, which will help to quickly come up with speed with how to perform Chabauty’s method in Magma.

Magma has a free, limited use online calculator here

<http://magma.maths.usyd.edu.au/calc/>,

and a thoroughly documented implementation of Chabauty’s method

<http://magma.maths.usyd.edu.au/magma/handbook/text/1533>.

Even better is to obtain a copy for your laptop, or ssh access to a departmental server with a copy of Magma. The Simons Foundation has graciously made Magma freely available to mathematicians working in the US

<http://magma.maths.usyd.edu.au/magma/ordering/>;

your department’s tech staff should be able to help you obtain a copy of Magma through this agreement.

Exercise 2.3. Take Smart’s list (from [Sma97]) of the 427 genus 2 curves with good reduction away from 2, and provably find all of the rational points on them. A temporary folder containing several references, and containing a subfolder titled “preparatory-Magma-exercise” with instructions for this exercise, is available at

<http://www.math.emory.edu/~dzb/AWS2020>.

As an entry point to some of the additional computational techniques one might need (such as étale descent), we recommend Poonen’s surveys [Poo96] and [Poo02].

3. THE UNIFORMITY CONJECTURE

The *uniformity conjecture* is one of the outstanding open conjectures in arithmetic and diophantine geometry. Initially, Mazur asked whether one can bound $\#X(K)$ purely in terms of the rank of the Jacobian of X (see [Maz00, Page 223] [Maz86, Page 234]). This was later promoted to the following stronger conjecture.

Conjecture 3.1 ([CHM97]). Let K be a number field and let $g \geq 2$ be an integer. There exists a constant $B_g(K)$ such that for every smooth curve X over K of genus g , the number $\#X(K)$ of K -rational points is at most $B_g(K)$.

The uniformity conjecture famously follows [CHM97, Theorem 1.1] from the Weak Lang conjecture (a higher dimension analogue of the Mordell conjecture), which is the following.

Conjecture 3.2 ([Lan74], 1.3; see also [Lan86]). Let X be a smooth proper variety of general type over a number field K . Then there exists a proper closed subscheme Z of X such that $X(K) = Z(K)$.

Alternatively, there are the following stronger pair of conjectures.

Conjecture 3.3 (Generic Uniform Boundedness [CHM97]). Let $g \geq 2$ be an integer. There exists a constant B_g such that for number field K , there exist only finitely many isomorphism classes of curves of genus g and over K such that $\#X(K) > B_g$.

This follows from the Strong Lang Conjecture.

Conjecture 3.4 ([Lan74], 1.3; see also [Lan86]). Let X be a smooth proper variety of general type over a number field K . Then there exists a proper closed subscheme Z of X such that for every finite extension $K \subset L$, the complement $X(L) - Z(L)$ is finite.

In [CHM97], one applies the Weak (or Strong) Lang Conjecture to symmetric powers of the universal curve $\mathcal{C} \rightarrow \overline{M}_{g,n}$. A major aspect of their proof is to show that large enough symmetric powers of \mathcal{C} are of general type (or at least dominate a variety of general type); this is a special case of their “correlation” theorem [CHM97, Theorem 1.2]. See the papers [Pac97, Pac99, Abr97, Abr95, Cap95, CHM95] for improvements, variants and a lot of additional discussion, and the slides

<http://www-math.mit.edu/~poonen/slides/uniformboundedness.pdf>

for a fairly recent discussion and some additional motivation.

3.5. Evidence and records. The following table (taken from [Cap95, Section 4] gives the best known lower bounds on the constant $B_g(\mathbb{Q})$.

g	2	3	4	5	10	45	g
$B_g(\mathbb{Q}) \geq$	642	112	126	132	192	781	$16(g+1)$

The record so far is due to Michael Stoll, who found (searching systematically through several families of curves constructed by Noam Elkies) the following:

$$y^2 = 82342800x^6 - 470135160x^5 + 52485681x^4 + 2396040466x^3 + 567207969x^2 - 985905640x + 247747600$$

It has at least 642 rational points, and rank at most 22. See

<http://www.mathe2.uni-bayreuth.de/stoll/recordcurve.html>

for a full list of the known points.

The families constructed by Elkies arise in the following way: he studied K3 surfaces of the form

$$y^2 = S(t, u, v)$$

with lots of rational lines, such that S restricted to such a line is a perfect square.

3.6. Chabauty–Coleman bounds. The proofs of Mordell due to Faltings, Vojta, and Bombieri [Fal97, Voj91, Bom90] give upper bounds on $\#X(K)$. These bounds tend to be astronomical, and are not explicit in their original proofs; moreover, it is unclear (to me) how they depend on X and K .

One application of Chabauty’s method is to give uniform bounds on small rank curves. Coleman’s original theorem is the following.

Theorem 3.7 (Coleman, [Col85]). *Let X/\mathbb{Q} be a curve of genus g and let $r = \text{rank}_{\mathbb{Z}} \text{Jac}_X(\mathbb{Q})$. Suppose $p > 2g$ is a prime of **good reduction**. Suppose $r < g$. Then*

$$\#X(\mathbb{Q}) \leq \#X(\mathbb{F}_p) + 2g - 2.$$

See [Col85, Lemma 3] or [MP12, Theorem 5.3] for a proof.

Various authors have worked to weaken Coleman’s hypotheses and to improve the bound; see Theorems 4.1, 5.1, 5.2, 6.1, and 6.2 below and the surrounding discussion.

3.8. A new hope. The recent work of Dimitrov, Gao and Habegger [DGH19, DGH20] give bounds on $\#X(K)$ which only depend on $g(X)$, $\deg K$, and $\text{rank Jac}_X(K)$. Combined with the conjectural boundedness of ranks of Jacobians of curves of fixed genus over a fixed number field (as predicted by [PPVW19] and [Poo18, Section 4.2]) this would prove uniformity. Their approach is in the spirit of Vojta’s original proof [Voj91], and relies on their recent other work on improved height bounds.

4. BAD REDUCTION

One avenue to improve on Coleman’s bound is to generalize the Chabauty framework and Coleman’s arguments to the case of bad reduction. We will discuss the advantage of working at bad primes and the difficulties that arise, starting with the work of Lorenzini–Tucker [LT02].

Coleman’s original bound (3.7) relies on an initial choice of a prime of good reduction. The first such prime could be arbitrarily large (e.g., consider a hyperelliptic curve $y^2 = f(x)$, and twist it by d , where d is the product of the first million primes; or, pick singular curves X_p over \mathbb{F}_p , for the first million primes p , and use the Chinese Remainder Theorem to construct a curve X over \mathbb{Q} that $X_{\mathbb{F}_p} \cong X_p$ for each such prime p). The problem with this is that the Hasse bound only provides that $\#X(\mathbb{F}_p) \leq 2g\sqrt{p} + p + 1$; so as p increases, Coleman’s bound becomes increasingly worse, and in particular is not uniform.

Whence the appeal of the following theorem of Lorenzini and Tucker.

Theorem 4.1 (Lorenzini, Tucker, [LT02], Corollary 1.11). *Suppose $p > 2g$ and let \mathcal{X} be a proper regular model of X over \mathbb{Z}_p . Suppose $r < g$. Then*

$$\#X(\mathbb{Q}) \leq \#\mathcal{X}_{\mathbb{F}_p}^{\text{sm}}(\mathbb{F}_p) + 2g - 2$$

where $\mathcal{X}_{\mathbb{F}_p}^{\text{sm}}$ is the smooth locus of the special fiber $\mathcal{X}_{\mathbb{F}_p}$.

Recall that a scheme X is **regular** if for every point $x \in X$, with corresponding maximal ideal \mathfrak{m} and residue field $k(x)$,

$$\dim_{k(x)} \mathfrak{m}/\mathfrak{m}^2 = \dim X.$$

If R is a DVR with uniformizer π and residue field k and $X \rightarrow \text{Spec } R$ is a relative curve, then a point $x \in X_k$ is regular if and only if the local equation at x is $yz = \pi$. (By “local equation” we mean the equation for the completion of the étale local ring $\mathcal{O}_{X,x}$.) See the examples in Subsection 4.2, and see [Sil94, Chapter IV] for a leisurely treatment.

The utility of the proper regular model \mathcal{X} of X is that the reduction map

$$r: \mathcal{X}(\mathbb{Q}) \rightarrow \mathcal{X}(\mathbb{F}_p)$$

takes values in the smooth locus $\mathcal{X}^{\text{sm}}(\mathbb{F}_p)$. In Chabauty’s method, one thus only needs to consider residue classes $r^{-1}(Q)$ of points $Q \in \mathcal{X}^{\text{sm}}(\mathbb{F}_p)$. Such residue classes are (p -adically analytically) isomorphic to discs; this makes the setup of Chabauty easier, and makes the “local analysis” much easier.

The “ $2g - 2$ ” term in Coleman’s bound (3.7) is derived from Riemann–Roch on $X_{\mathbb{F}_p}$, and is the rationale for the “good reduction” hypothesis. Lorenzini and Tucker recover the $2g - 2$ term via Riemann–Roch on $X_{\mathbb{Q}_p}$ and a more involved p -adic analytic argument. A later, alternative proof [MP12, Theorem A.5] instead recovers the $2g - 2$ term via arithmetic intersection theory on \mathcal{X} and adjunction.


The drawback is that $\mathcal{X}_{\mathbb{F}_p}$ could contain an arbitrarily long chain of \mathbb{P}^1 ’s. (For example, if X is an elliptic curve with semistable reduction at p and $v_p(j(X)) = -n$, then $\mathcal{X}_{\mathbb{F}_p}$ is an n -gon of \mathbb{P}^1 ’s.) Once again, the size of $\mathcal{X}^{\text{sm}}(\mathbb{F}_p)$ could be arbitrarily large, giving non-uniform bounds.

Stoll [Sto19] had the bold idea to work with a *non-regular, minimal* model: one contracts each chain of \mathbb{P}^1 ’s into a single node. Such a model is no longer regular, but the number of components is bounded, and the genus of each component is bounded (exercise: verify this). Since the model is no longer regular, rational points no longer reduce to smooth points (exercise: give an example), and might reduce to a node. The residue class of a node is now an annulus (explain in an example). This creates multiple problems: an annulus admits “monodromy” and integrals no longer admit a unique analytic continuation, and local expansions are now Laurent, rather than power, series. See Theorem 6.1 below and the surrounding discussion.

4.2. A few examples.

Example 4.3 (A regular model). The relative curve


$$\begin{aligned} y^2 &= (x(x-1)(x-2))^3 - 5 \\ &= (x(x-1)(x-2))^3 \pmod{5}. \end{aligned}$$




is regular at the point $(0, 0)$. The local equation at $(0, 0)$ analytically looks like $xy = 5$. Once can see by elementary number theory that no rational point can reduce to $(0, 0)$.

Example 4.4 (Resolving a semistable, but not regular, model).


$$\begin{aligned} y^2 &= (x(x-1)(x-2))^3 - 5^4 \\ &= (x(x-1)(x-2))^3 \pmod{5} \end{aligned}$$



Now, the local equation at $(0,0)$ looks like $xy = 5^4$, and $(0, 5^2)$ reduces to $(0, 0)$. Blowing up along the ideal $(x, y, 5)$ gives



and the local equations now look like $xy = 5^3$ and $xy = 5$. One of the 2 points is still not regular. After 2 more blowups we get



and now all of the local equations look like $xy = 5$, giving a regular model.

5. RANK FAVORABLE BOUNDS

Lorenzini and Tucker [LT02] ask if one can refine Coleman’s bound (Theorem 3.7) when the rank r is small (i.e., $r \leq g - 2$). This was subsequently answered by Stoll.

Theorem 5.1 (Stoll, [Sto06], Corollary 6.7). *With the hypothesis of Theorem 3.7,*

$$\#X(\mathbb{Q}) \leq \#X(\mathbb{F}_p) + 2r.$$

The space of differentials “suitable” for Chabauty’s method has dimension at least $g - r$. When $r < g - 1$, there are thus more “inputs” to Chabauty’s method and this extra flexibility can be exploited to improve the p -adic analysis. Indeed, Stoll’s idea is: instead of using a single integral, to tailor the choice of integral to each residue class. The additional “global geometric input” is the (classical) notion of “rank of a divisor”, and after translating his setup into this language, improved bounds follow from Clifford’s Theorem.

The application of Clifford’s theorem is also the source of the “good reduction” hypothesis. Using ideas from the “discrete case” of tropical geometry (in particular “chip-firing”), Eric Katz and I generalized Stoll’s theorem to arbitrary reduction types.

Theorem 5.2 (Katz, Zureick-Brown, [KZB13]). *Let X/\mathbb{Q} be a curve of genus g and let $r = \text{rank Jac}_X(\mathbb{Q})$. Suppose $p > 2r + 2$ is a prime, that $r < g$, and let \mathcal{X} be a proper regular*

model of X over \mathbb{Z}_p . Then

$$\#X(\mathbb{Q}) \leq \#\mathcal{X}_{\mathbb{F}_p}^{sm}(\mathbb{F}_p) + 2r.$$

Unlike Lorenzini and Tucker’s generalization of Coleman’s theorem, where they replace Coleman’s use of Riemann–Roch on $X_{\mathbb{F}_p}$ with Riemann–Roch on $X_{\mathbb{Q}_p}$, it does not seem possible to replace Stoll’s use of Clifford’s theorem on $X_{\mathbb{F}_p}$ with Clifford’s theorem on $X_{\mathbb{Q}_p}$. Matt Baker suggested that it might be possible to generalize Stoll’s theorem to curves with bad, *totally degenerate* reduction (i.e., $X_{\mathbb{F}_p}$ is a union of rational curves meeting transversely) using ideas from tropical geometry (see the recent survey [BJ15] on tropical geometry and applications), in particular the notion of “chip firing”, Baker’s combinatorial definition of rank, and Baker–Norine’s [BN07] combinatorial Riemann–Roch and Clifford theorems. Baker was correct, and in fact an enrichment of his theory led to the following common generalization of Stoll’s and Lorenzini and Tucker’s theorems.

Baker’s recent work [Bak08] clarifies the relationship between *linear systems* on curves and on finite graphs. Highlights include a semicontinuity theorem for ranks of linear systems (as one passes from the curve to its dual graph), and graph theoretic analogues of Riemann–Roch and Clifford’s theorem. Baker’s theory works best with totally degenerate curves (i.e. each component is a \mathbb{P}^1). Theorem 5.2 requires an enrichment of Baker’s theory if the irreducible components of the reduction have higher genus.

5.3. Stoll’s proof of Theorem 5.1. Let $p > 2$. Denote by V_{chab} the vector space of all $\omega \in H^0(X_{\mathbb{Q}_p}, \Omega_{X_{\mathbb{Q}_p}/\mathbb{Q}_p}^1)$ such that $\int_{P_1}^{P_2} \omega = 0$ for all $P_1, P_2 \in X(\mathbb{Q})$. Then $\dim V_{\text{chab}} \geq g - r$. For each $\omega \in V_{\text{chab}}$, scale ω by a power of p so that the reduction $\tilde{\omega}$ of ω is non zero, and denote by \tilde{V}_{chab} the set of all such reductions; we note that $\dim_{\mathbb{F}_p} \tilde{V}_{\text{chab}} = \dim_{\mathbb{Q}_p} V_{\text{chab}}$.

For each $Q \in X(\mathbb{F}_p)$ and $\omega \in \tilde{V}_{\text{chab}}$, set

$$n_Q(\omega) := \deg(\text{div } \omega|_{]Q[}) \text{ and } n_Q := \min_{\omega \in \tilde{V}_{\text{chab}}} n_Q(\omega)$$

(where we recall that $]Q[$ denotes the *tube* or *residue class* of Q , that is, the set of all points of $X(\mathbb{Q}_p)$ which reduce to Q). Since div is compatible with reduction mod p , $n_Q(\omega)$ is equal to the valuation $v_Q(\tilde{\omega})$ (i.e., the order of vanishing of $\tilde{\omega}$ at Q).

By the “ p -adic Rolle’s theorem”, the number of zeroes of $\int \omega$ in $]Q[$ is at most $1 + n_Q(\omega)$, so

$$X(\mathbb{Q}) \leq \sum_{Q \in X(\mathbb{F}_p)} (1 + n_Q) = \sum_{Q \in X(\mathbb{F}_p)} 1 + \sum_{Q \in X(\mathbb{F}_p)} n_Q = X(\mathbb{F}_p) + \deg(D_{\text{chab}}),$$

where we define D_{chab} to be the divisor

$$D_{\text{chab}} := \sum_{Q \in X(\mathbb{F}_p)} n_Q Q \in \text{Div } X_{\mathbb{F}_p}.$$

By Riemann–Roch, $\deg D_{\text{chab}} \leq 2g - 2$, recovering the bound

$$X(\mathbb{Q}) \leq X(\mathbb{F}_p) + 2g - 2.$$

We claim that, in fact, $\deg D \leq 2r$, which suffices to prove Theorem 5.1. (When $r = g - 1$, $2r = 2g - 2$.) Stoll’s main observation is that

$$\tilde{V}_{\text{chab}} \subset H^0(X_{\mathbb{F}_p}, \Omega_{X_{\mathbb{F}_p}}^1(-D_{\text{chab}})), \tag{5.3.1}$$

and in particular,

$$\dim H^0(X_{\mathbb{F}_p}, \Omega_{X_{\mathbb{F}_p}}^1(-D_{\text{chab}})) \geq \dim \tilde{V}_{\text{chab}} \geq g - r. \quad (5.3.2)$$

To justify Equation 5.3.1, given an effective divisor $E = \sum n_P P$ and a line bundle \mathcal{L} on a curve X , recall that $H^0(X, \mathcal{L}(-E))$ is the subspace of sections of $H^0(X, \mathcal{L})$ that have at least a zero of order n_P at P . A differential $\tilde{\omega} \in \tilde{V}_{\text{chab}}$ thus satisfies $v_P(\omega) \geq n_P$ by definition of n_P !

On the other hand, Clifford's Theorem [Har77, Theorem IV.5.4] implies that

$$\dim H^0(X_{\mathbb{F}_p}, \Omega_{X_{\mathbb{F}_p}}^1(-D_{\text{chab}})) \leq \frac{1}{2} \deg \left(\Omega_{X_{\mathbb{F}_p}}^1(-D_{\text{chab}}) \right) + 1. \quad (5.3.3)$$

Combining equations 5.3.2 and 5.3.3 gives

$$g - r \leq \frac{1}{2} \deg \left(\Omega_{X_{\mathbb{F}_p}/\mathbb{Q}_p}^1(-D_{\text{chab}}) \right) + 1 = g - 1 - \frac{1}{2} \deg D_{\text{chab}} + 1$$

and simplifying gives

$$\deg D_{\text{chab}} \leq 2r.$$

To justify Equation 5.3.3, we switch to the language of divisors. Recall that a divisor is *special* if $\dim H^0(X, K - D) > 0$, where K is a canonical divisor. (Equivalently, D is special if and only if it is a subdivisor of some canonical divisor.) For context: Riemann–Roch reads:

$$H^0(X, D) = \dim H^0(X, K - D) + \deg D + 1 - g.$$

This gives a formula for $H^0(X, D)$ when the degree of D is large; indeed when $\deg D > \deg K = 2g - 2$, $\deg K - D < 0$, therefore $\dim H^0(X, K - D) = 0$ and $H^0(X, D) = \deg D + 1 - g$. At the other extreme: if $\deg D \leq 2g - 2$, then it is still possible that $\dim H^0(X, K - D) = 0$, in which case, again, $H^0(X, D) = \deg D + 1 - g$. If D is special, i.e., $\dim H^0(X, K - D) > 0$, then by Riemann–Roch, $H^0(X, D) < \deg D + 1 - g$; in this case, *Clifford's Theorem* gives the much stronger bound

$$H^0(X, D) \leq \frac{1}{2} (\deg D) + 1.$$

Let $K = \text{div } \tilde{\omega}$. Then, by definition of n_Q , D_{chab} is a subdivisor of the canonical divisor K (since $v_Q(D_{\text{chab}}) := v_Q(\tilde{\omega}) = n_Q(\omega) \geq n_Q$); in particular, D_{chab} is special.

5.4. The rank of a divisor. In the proof of Stoll's theorem, we implicitly used the notion of *rank of a line bundle* (or divisor). One can simply define the rank $r(\mathcal{L})$ of a divisor $\mathcal{L} \in \text{Pic}(X)$ to be

$$r(\mathcal{L}) := \dim H^0(X, \mathcal{L}) - 1.$$

This has the following alternative interpretation over an algebraically closed field: $r(\mathcal{L})$ is the largest number of independent and generic “vanishing conditions” one can impose on sections of \mathcal{L} . Recall that for a closed point P , $H^0(X, \mathcal{L}(-P))$ is the subspace of sections of $H^0(X, \mathcal{L})$ that have at least a simple zero at the point P . More generally, for an effective divisor $E = \sum n_P P$, $H^0(X, \mathcal{L}(-E))$ is the subspace of sections of $H^0(X, \mathcal{L})$ that have at least a zero of order n_P at P . By [Har77, Proof of Theorem IV.1.3],

$$\dim H^0(X, \mathcal{L}(-E)) \geq \dim H^0(X, \mathcal{L}) - 1,$$

and in particular,

$$\dim H^0(X, \mathcal{L}(-E)) \geq \dim H^0(X, \mathcal{L}) - \deg E. \quad (5.4.1)$$

Similarly, one can define the rank $r(D)$ of a divisor $D \in \text{Div}(X)$ to be

$$r(D) := \dim H^0(X, D) - 1.$$

This has the following alternative interpretation over an algebraically closed field: $r(D)$ is the largest number of points of $X_{\bar{k}}$ (allowing for multiplicity) one can remove from D before D is no longer equivalent to some effective divisor. Equivalently, the rank is the largest number of points (allowing for multiplicity) that you can demand occurs as a subdivisor of some effective divisor D' equivalent to D .

More formally, we make the following definitions.

Definition 5.5. Let $D \in \text{Div } X$ be a divisor. The **linear system** associated to D is the collection $|D|$ of effective divisors linearly equivalent to D . We define the **rank** $r(D)$ of D to be -1 if $|D|$ is empty (i.e., if D is not equivalent to an effective divisor). Otherwise, we define

$$r(D) := \max\{n \in \mathbb{Z}_{\geq 0} : |D - E| \neq \emptyset, \forall E \in \text{Div}_{\geq 0}^n(X_{\bar{k}})\},$$

where $\text{Div}_{\geq 0}^n(X_{\bar{k}})$ is the subset of $\text{Div}(X_{\bar{k}})$ of effective divisors of degree n .

The linear system $|D|$ is naturally isomorphic to the projective space $\text{Proj } H^0(X, D) \cong \mathbb{P}^r - 1$, where $r = \dim H^0(X, D)$.

By Equation 5.4.1,

$$r(D) \geq \dim H^0(X, D) - 1.$$

The converse follows from the observation (exercise!) that if $r(D) = n$, then there exists $P \in X_{\bar{k}}$ such that $r(D - P) = n - 1$. Note that one must take \bar{k} in the definition of rank. Indeed, consider a non hyperelliptic curve X with $X(k) = \emptyset$, but $X(k') \neq \emptyset$ for some quadratic extension k' of k . Then for $P \in X(k')$ with conjugate point Q , $D := P + Q \in \text{Div}^2 X$. Then $r(D) = 1$; but, $\text{Div}_{\geq 0}^1 X$ is empty, so

$$\max\{n \in \mathbb{Z}_{\geq 0} : |D - E| \neq \emptyset, \forall E \in \text{Div}_{\geq 0}^n(X)\} = 1$$

5.6. Chip firing and the rank of a divisor on a graph. I recommend taking a quick look at Matt Baker's short expository article available at

<http://people.math.gatech.edu/~mbaker/pdf/g4g9.pdf>.

For a short selection of other references: the papers [Bak08, BN07] by Baker and collaborators are my preferred starting point; [BJ15] is also a great survey.

Let Γ be a connected graph, with vertex set $V(\Gamma)$ and edge set $E(\Gamma)$. We define $\text{Div } \Gamma$ to be the set of maps from the vertices of $V(\Gamma)$ to \mathbb{Z} ; this is isomorphic to the free group $\mathbb{Z}[V(\Gamma)]$ generated by the set of vertices of Γ , and will sometimes write $D = \sum_v n_v(v)$ for the function that has value n_v at v . The **degree** of D is $\deg D = \sum_v D(v)$. We'll refer to an element $D \in \text{Div } \Gamma$ as a **divisor** or **configuration**, and will typically represent them visually as in Figure 5.6, and refer to $D(v)$ as the “number of chips” or “dollars” at the vertex v .

The goal of the “dollar game” or “chip firing” is to get a divisor out of debt. We say that a divisor D is **effective**, and write $D \geq 0$, if $D(v) \geq 0$ for all vertices v of Γ .

One formalizes lending and borrowing as follows: given $f \in \text{Div } \Gamma$, we define the principal divisor associated to f to be

$$\text{div } f = \sum_{v \in \Gamma} D(v) \cdot \left(-(\deg v)(v) + \sum_{w \neq v} \#\{\text{edges between } w \text{ and } v\}(w) \right).$$




FIGURE 1. The effect of firing once at v_4

In particular, $\text{div } \delta_v$ is

$$-(\deg v)(v) + \sum_{w \neq v} \#\{\text{edges between } w \text{ and } v\}(w)$$

which has the effect of the vertex v “lending” one chip to each adjacent vertex (see Figure 5.6). We say that two divisors D and D' are **equivalent** if there is a sequence of lends and borrows which transforms D into D' , and we define the **Jacobian** or **Picard group** $\text{Pic } \Gamma$ to be the abelian group of equivalence classes of divisors on Γ . more formally, there is an exact sequence

$$0 \rightarrow Z \rightarrow \text{Div } \Gamma \xrightarrow{\text{div}} \text{Div } \Gamma \rightarrow \text{Pic } \Gamma \rightarrow 0,$$

where the first map sends 1 to the function $\sum_v \delta_v$ (i.e., every vertex lends, which has no effect).

The vector space $H^0(\Gamma, D)$ doesn't make sense for a graph. Baker's insight from [Bak08] is that the “practical” definition of rank (Definition 5.5) *does* generalize nicely to divisors on graphs.

Definition 5.7. Let $D \in \text{Div } \Gamma$ be a divisor. The **linear system** associated to D is the collection $|D|$ of effective divisors linearly equivalent to D . We define the **rank** $r(D)$ of D to be -1 if $|D|$ is empty (i.e., if D is not equivalent to an effective divisor). Otherwise, we define

$$r(D) := \max\{n \in \mathbb{Z}_{\geq 0} : |D - E| \neq \emptyset, \forall E \in \text{Div}_{\geq 0}^n(\Gamma)\},$$

where $\text{Div}_{\geq 0}^n(X_{\bar{k}})$ is the subset of $\text{Div}(X_{\bar{k}})$ of effective divisors of degree n .

Equivalently, the rank is the largest number of points (allowing for multiplicity) that you can demand occurs as a subdivisor of some effective divisor D' equivalent to D . In other words, the rank of a divisor D is the “amount of damage” necessary to make the chip firing game unwinnable.

Definition 5.8. Let Γ be a graph. We define the **canonical divisor** on K to be the divisor

$$K_{\Gamma} := \sum_{v \in V(\Gamma)} (\deg v - 2)(v).$$

The **genus** $g(\Gamma)$ (alternatively **first Betti number** $h^1(\Gamma)$) of Γ is the number of edges minus the number of vertices; note that $\deg K_{\Gamma} = 2g(\Gamma) - 2$. We say that a divisor D is **special** if D is effective and $|K - D|$ is non empty.

This definition is motivated by adjunction.

Theorem 5.9 ([BN07], Theorem 1.12 and Corollary 3.5). *Let $D \in \text{Div } \Gamma$. Then the following are true.*

- (1) *Riemann–Roch: $r(D) - r(K - D) = \deg D + 1 - g$.*
- (2) *Clifford’s Theorem: if D is special, then $r(D) \leq \frac{1}{2} \deg D$.*

Corollary 5.10 ([BN07], Theorem 1.9). *The chip firing game is winnable for the configuration D if $\deg D \geq g$.*

Remark 5.11. It is unknown whether one can deduce these from the analogous theorems from the geometry of curves.

5.12. Semicontinuity of specialization. Let R be a complete discrete valuation ring with maximal ideal π , residue field k , and fraction field K . Denote by η the generic point of $\text{Spec } R$, and by b the closed point. (Most of what we say below works just as well if we replace $\text{Spec } R$ by an integral scheme B .) Let $C \rightarrow B$ be a relative curve of genus g (i.e., a smooth proper morphism such that for every $x \in \text{Spec } R$, the fiber C_x is a smooth proper curve of genus g over the residue field $k(x)$).

Let $D = \sum n_P P \in \text{Div } C_\eta$. The dimension of C is 2, and we can extend D to a divisor \mathcal{D} on C by taking the closure of its support; in other words, $\mathcal{D} := \sum n_P \bar{P} \in \text{Div } C_\eta$, where \bar{P} is the closure of P . Intersecting \mathcal{D} with the special fiber C_b thus gives a specialization map

$$\text{sp}: \text{Div } C_\eta \rightarrow \text{Div } C_b.$$

Proposition 5.13. *Let $D \in \text{Div } C_\eta$. Then $r(\text{sp}(D)) \geq r(D)$.*

The inequality can certainly be strict. Indeed, consider \mathcal{C} with hyperelliptic special fiber and non-hyperelliptic generic fiber, and let $D = P + Q$ where $P, Q \in \mathcal{C}(K)$ are points whose reductions are hyperelliptic conjugate. Then $r(D) = 1$ but $r(\text{sp}(D)) = 2$.

More generally, if L is a line bundle on C_η , there is a unique (up to isomorphism) extension of L to a line bundle \mathcal{L} on C (i.e., a line bundle \mathcal{L} on C such that \mathcal{L}_η is isomorphic to L . (Indeed, let $s \in L(U)$ be a section over some non empty open set $U \subset C_\eta$; then $\mathcal{L} = \mathcal{O}_C(\bar{\cdot} \div s)$ extends L .)

There is thus an analogous specialization map

$$\text{sp}: \text{Pic } C_\eta \rightarrow \text{Pic } C_b,$$

and (since $r(D) = r(\mathcal{O}(D))$), Proposition 5.13 equivalently implies that $r(\text{sp}(L)) \geq r(L)$.

Proposition 5.13 is a special case of Semicontinuity of Cohomology [Har77, Theorem III.12.8] (taking $i = 0$ and $\mathcal{F} = \mathcal{O}_C(D)$). Unsurprisingly, this is overkill; we sketch a direct proof that will generalize to the “discrete” case.

Proof of Proposition 5.13. First, note that a section $s \in H^0(C_\eta, D)$ can be scaled by a power of the uniformizer π to a section of $H^0(C, \mathcal{O}(\bar{D}))$. There are a few ways to see this: if we think of s as a function

Conversely, Since $C_\eta \subset C$ is open, and since the vanishing locus of a section of a line bundle is closed, the map

$$H^0(C, \mathcal{O}(\bar{D})) \rightarrow H^0(C_\eta, D)$$

is injective. In particular,

$$\text{rank}_R H^0(C, \mathcal{O}(\bar{D})) = \text{rank}_K H^0(C_\eta, D).$$



FIGURE 2. A curve and its dual graph.

Alternatively, it follows directly from flatness that

$$H^0(C, \mathcal{O}(\overline{D})) \otimes_R K \cong H^0(C_\eta, D).$$

One can then show that the dimension of the reduction map

$$H^0(C, \mathcal{O}(\overline{D})) \rightarrow H^0(C_b, \mathcal{O}(D_b))$$

is $\dim H^0(C_\eta, D)$, so in particular

$$\dim H^0(C_b, \mathcal{O}(D_b)) \geq \dim H^0(C_\eta, D).$$

□

5.14. Rank favorable bounds for curves with totally degenerate reduction. One can associate to such a singular curve with transverse crossings its dual graph as in Figure 1. Component curves become nodes, and intersections correspond to edges.

In this lectures, I'll discuss the special case of a “Mumford curve” (i.e., a curve with “totally degenerate” reduction, in that the reduction is a collection of \mathbb{P}^1 's meeting transversely, and in particular represents an isolated point on the moduli space of curves); for such curves, one only needs Baker's original notion of rank (which we call “numerical rank”); for a discussion of the “abelian rank”, and a detailed proof in this case, see our paper [KZB13]. The main point is that there is also a specialization map for the numerical (i.e., “chip firing”) rank, and once one sets things up properly, the proof is similar to Stoll's proof.

5.15. Refined ranks. In a certain sense, the numerical rank only “sees” the component group of the Néron model. The enriched notion of abelian rank from [KZB13] and [AB15] sees the abelian part of the Néron model. One can ask if there is a corresponding notion of “toric” or “unipotent” rank. In [KZB13, Subsection 3.3], we define a “toric” rank, and in [KZB13, Example 5.5] demonstrate that these ranks differ; we have yet to find a useful application.

6. TROPICAL GEOMETRY, BERKOVICH SPACES, AND CHABAUTY

A recent breakthrough [Sto19] fully removed, in the special case of hyperelliptic curves, the dependence on a *regular* model and derived a *uniform* bound on $\#X(\mathbb{Q})$ for small ($r \leq g-3$) rank curves.

Theorem 6.1 (Stoll, [Sto19]). *Let X be a hyperelliptic curve of genus g and let $r = \text{rank}_{\mathbb{Z}} \text{Jac}_X(\mathbb{Q})$. Suppose $r \leq g-3$. Then*

$$\#X(\mathbb{Q}) \leq 8(r+4)(g-1) + \max\{1, 4r\} \cdot g.$$

A main ingredient in Stoll’s proof is to understand the discrepancy between the different flavors of integration. Eric Katz noticed that this discrepancy “factored through the tropicalization of the torus part of the Berkovich uniformization of X ”. After a thorough reinterpretation of the method of Chabauty and Coleman via Berkovich spaces, and harnessing the full catalogue of tropical and non-Archimedean analytic tools, we were able to improve Stoll’s result to *arbitrary* curves of small rank.

Theorem 6.2 (Katz–Rabinoff–Zureick-Brown, [KRZB]). *Let X be **any** curve of genus g and let $r = \text{rank}_{\mathbb{Z}} \text{Jac}_X(\mathbb{Q})$. Suppose $r \leq g - 3$. Then*

$$\#X(\mathbb{Q}) \leq 84g^2 - 98g + 28.$$

For more details, see our survey [KRZB16].

REFERENCES

- [AB15] Omid Amini and Matthew Baker, *Linear series on metrized complexes of algebraic curves*, Math. Ann. **362** (2015), no. 1-2, 55–106. MR3343870 ↑16
- [Abr95] Dan Abramovich, *Uniformité des points rationnels des courbes algébriques sur les extensions quadratiques et cubiques*, C. R. Acad. Sci. Paris Sér. I Math. **321** (1995), no. 6, 755–758. MR1354720 ↑7
- [Abr97] ———, *Uniformity of stably integral points on elliptic curves*, Invent. Math. **127** (1997), no. 2, 307–317. MR1427620 ↑7
- [Bak08] Matthew Baker, *Specialization of linear systems from curves to graphs*, Algebra Number Theory **2** (2008), no. 6, 613–653. With an appendix by Brian Conrad. MR2448666 (2010a:14012) ↑3, 4, 11, 13, 14
- [BD19] Jennifer S. Balakrishnan and Netan Dogra, *An effective Chabauty-Kim theorem*, Compos. Math. **155** (2019), no. 6, 1057–1075. MR3949926 ↑5
- [BJ15] Matthew Baker and David Jensen, *Degeneration of linear series from the tropical point of view and applications*, arXiv preprint arXiv:1504.05544 (2015). ↑3, 11, 13
- [BN07] M. Baker and S. Norine, *Riemann-Roch and Abel-Jacobi theory on a finite graph*, Adv. Math. **215** (2007), no. 2, 766–788. MR2355607 (2008m:05167) ↑11, 13, 14
- [Bom90] E. Bombieri, *The Mordell conjecture revisited*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **17** (1990), no. 4, 615–640. MR1093712 (92a:11072) ↑2, 7
- [BPR13] Matthew Baker, Sam Payne, and Joseph Rabinoff, *On the structure of nonarchimedean analytic curves*, Tropical and Non-Archimedean Geometry, 2013, pp. 93–121. ↑4
- [BPS16] Nils Bruin, Bjorn Poonen, and Michael Stoll, *Generalized explicit descent and its application to curves of genus 3*, Forum Math. Sigma **4** (2016), e6, 80. MR3482281 ↑2
- [Cap95] L. Caporaso, *Counting rational points on algebraic curves*, Rend. Sem. Mat. Univ. Politec. Torino **53** (1995), no. 3, 223–229. Number theory, I (Rome, 1995). MR1452380 (98i:11041) ↑7
- [Cas83] J. W. S. Cassels, *The Mordell-Weil group of curves of genus 2*, Arithmetic and geometry, Vol. I, 1983, pp. 27–60. MR717589 ↑2
- [Cas89] ———, *Arithmetic of curves of genus 2*, Number theory and applications (Banff, AB, 1988), 1989, pp. 27–35. MR1123068 ↑2
- [Cha41] C. Chabauty, *Sur les points rationnels des courbes algébriques de genre supérieur à l’unité*, C. R. Acad. Sci. Paris **212** (1941), 882–885. ↑2, 5
- [CHM95] Lucia Caporaso, Joe Harris, and Barry Mazur, *How many rational points can a curve have?*, The moduli space of curves (Texel Island, 1994), 1995, pp. 13–31. MR1363052 (97d:11099) ↑7
- [CHM97] ———, *Uniformity of rational points*, J. Amer. Math. Soc. **10** (1997), no. 1, 1–35. MR1325796 (97d:14033) ↑6, 7
- [Col85] R. F. Coleman, *Effective Chabauty*, Duke Math. J. **52** (1985), no. 3, 765–770. ↑2, 5, 7
- [DGH19] Vesselin Dimitrov, Ziyang Gao, and Philipp Habegger, *Uniform bound for the number of rational points on a pencil of curves*, arXiv preprint arXiv:1904.07268 (2019). ↑8

- [DGH20] ———, *Uniformity in mordell-lang for curves*, arXiv preprint arXiv:2001.10276 (2020). [↑8](#)
- [Fal86] Gerd Faltings, *Finiteness theorems for abelian varieties over number fields* (1986), 9–27. Translated from the German original [Invent. Math. **73** (1983), no. 3, 349–366; *ibid.* **75** (1984), no. 2, 381; MR 85g:11026ab] by Edward Shipz. MRFaltings:bookArithmeticGeometry [↑2](#)
- [Fal97] ———, *The determinant of cohomology in etale topology*, Arithmetic geometry (Cortona, 1994), 1997, pp. 157–168. MRFaltings:bookArithmeticGeometry (98k:14023) [↑7](#)
- [Fly90] Eugene Victor Flynn, *The Jacobian and formal group of a curve of genus 2 over an arbitrary ground field*, Math. Proc. Cambridge Philos. Soc. **107** (1990), no. 3, 425–441. MR1041476 [↑2](#)
- [GG93] Daniel M. Gordon and David Grant, *Computing the Mordell-Weil rank of Jacobians of curves of genus two*, Trans. Amer. Math. Soc. **337** (1993), no. 2, 807–824. MR1094558 [↑2](#)
- [Har77] R. Hartshorne, *Algebraic geometry*, Springer-Verlag, New York-Heidelberg, 1977. Graduate Texts in Mathematics, No. 52. MR0463157 (57 #3116) [↑4](#), 5, 12, 15
- [KRZB16] Eric Katz, Joseph Rabinoff, and David Zureick-Brown, *Diophantine and tropical geometry, and uniformity of rational points on curves*, Survey article for the 2015 Summer Research Institute on Algebraic Geometry Proceedings (in review) (2016). [↑3](#), 17
- [KRZB] ———, *Uniform bounds for the number of rational points on curves of small mordell-weil rank*, to appear in Duke Mathematical Journal. [↑3](#), 4, 5, 17
- [KZB13] Eric Katz and David Zureick-Brown, *The Chabauty-Coleman bound at a prime of bad reduction and Clifford bounds for geometric rank functions*, Compos. Math. **149** (2013), no. 11, 1818–1838. MR3133294 [↑3](#), 10, 16
- [Lan74] Serge Lang, *Higher dimensional diophantine problems*, Bull. Amer. Math. Soc. **80** (1974), 779–787. MR360464 [↑6](#), 7
- [Lan86] ———, *Hyperbolic and Diophantine analysis*, Bull. Amer. Math. Soc. (N.S.) **14** (1986), no. 2, 159–205. MR828820 [↑6](#), 7
- [LT02] Dino Lorenzini and Thomas J. Tucker, *Thue equations and the method of Chabauty-Coleman*, Invent. Math. **148** (2002), no. 1, 47–77. [↑3](#), 8, 10
- [Maz00] Barry Mazur, *Abelian varieties and the Mordell-Lang conjecture*, Model theory, algebra, and geometry, 2000, pp. 199–227. MR1773708 (2001e:11061) [↑6](#)
- [Maz86] B. Mazur, *Arithmetic on curves*, Bull. Amer. Math. Soc. (N.S.) **14** (1986), no. 2, 207–259. MR828821 (88e:11050) [↑6](#)
- [McC94] William G. McCallum, *On the method of Coleman and Chabauty*, Math. Ann. **299** (1994), no. 3, 565–596. MR1282232 [↑2](#)
- [MP12] William McCallum and Bjorn Poonen, *The method of Chabauty and Coleman*, Explicit methods in number theory, 2012, pp. 99–117. MR3098132 [↑3](#), 4, 5, 7, 9
- [Pac97] Patricia L. Pacelli, *Uniform boundedness for rational points*, Duke Math. J. **88** (1997), no. 1, 77–102. MR1448017 [↑7](#)
- [Pac99] ———, *Uniform bounds for stably integral points on elliptic curves*, Proc. Amer. Math. Soc. **127** (1999), no. 9, 2535–2546. MR1676288 [↑7](#)
- [Poo02] Bjorn Poonen, *Computing rational points on curves*, Number theory for the millennium, III (Urbana, IL, 2000), 2002, pp. 149–172. [↑6](#)
- [Poo18] ———, *Heuristics for the arithmetic of elliptic curves*, Proceedings of the International Congress of Mathematicians—Rio de Janeiro 2018. Vol. II. Invited lectures, 2018, pp. 399–414. MR3966772 [↑8](#)
- [Poo96] ———, *Computational aspects of curves of genus at least 2*, Algorithmic number theory (Talence, 1996), 1996, pp. 283–306. MR1446520 (98c:11059) [↑6](#)
- [PPVW19] Jennifer Park, Bjorn Poonen, John Voight, and Melanie Matchett Wood, *A heuristic for boundedness of ranks of elliptic curves*, J. Eur. Math. Soc. (JEMS) **21** (2019), no. 9, 2859–2903. MR3985613 [↑8](#)
- [PS14] Bjorn Poonen and Michael Stoll, *Most odd degree hyperelliptic curves have only one rational point*, Ann. of Math. (2) **180** (2014), no. 3, 1137–1166. MR3245014 [↑2](#)
- [PSS07] Bjorn Poonen, Edward F. Schaefer, and Michael Stoll, *Twists of $X(7)$ and primitive solutions to $x^2 + y^3 = z^7$* , Duke Math. J. **137** (2007), no. 1, 103–158. MR2309145 (2008i:11085) [↑2](#)

- [RZB15] Jeremy Rouse and David Zureick-Brown, *Elliptic curves over \mathbb{Q} and 2-adic images of Galois representations*, Research in Number Theory, Accepted (2015). [↑2](#)
- [Sil94] J. H. Silverman, *Advanced topics in the arithmetic of elliptic curves*, Graduate Texts in Mathematics, vol. 151, Springer-Verlag, New York, 1994. [↑8](#)
- [Sko34] Thoralf Skolem, *Ein verfahren zur behandlung gewisser exponentialer gleichungen und diophantischer gleichungen*, C. r **8** (1934), 163–188. [↑2](#)
- [Sma97] N. P. Smart, *S-unit equations, binary forms and curves of genus 2*, Proc. London Math. Soc. (3) **75** (1997), no. 2, 271–307. MR1455857 [↑6](#)
- [Sto06] Michael Stoll, *Independence of rational points on twists of a given curve*, Compos. Math. **142** (2006), no. 5, 1201–1214. MR2264661 (2007m:14025) [↑3](#), [4](#), [10](#)
- [Sto19] ———, *Uniform bounds for the number of rational points on hyperelliptic curves of small Mordell-Weil rank*, J. Eur. Math. Soc. (JEMS) **21** (2019), no. 3, 923–956. MR3908770 [↑3](#), [4](#), [9](#), [16](#)
- [Voj91] P. Vojta, *Siegel’s theorem in the compact case*, Ann. of Math. (2) **133** (1991), no. 3, 509–548. MR1109352 (93d:11065) [↑2](#), [7](#), [8](#)

DEPT. OF MATHEMATICS, EMORY UNIVERSITY, ATLANTA, GA 30322 USA
E-mail address: `dzbo@mathcs.emory.edu`