

$$cl_{\mathbb{Q}} : CH^m(X) \otimes \mathbb{Q} \longrightarrow H^{2m}(X_{\mathbb{Q}}, \mathbb{Q}(m))$$

$$/\mathbb{C} : cl : CH^m(X)_{\mathbb{C}} \longrightarrow H^{2m}(X, \mathbb{C}(m))$$

Conj: should be  $\hookrightarrow$  on  $CH^i_{(0)}$

$\leadsto$   $CH^i_{(0)}$  are  $\mathbb{Q}$ -v. space of dim  $< \infty$

For  $CH^1_{(1)}$  and  $CH^2_{(1)}$  we see :

- in gen'l not of finite type
- depends on  $k$

Still : "manageable" because points of some alg. var.

Thm (Soulé, Künnemann)

If  $k \subseteq \overline{\mathbb{F}_p}$  then  $CH_{(s)}^m(X) = 0$   
for all  $s \neq 0$

$\subseteq CH^m(X)_{\mathbb{Q}}$  2

Idea of pf

• reduction to  $k = \mathbb{F}_q$ ,

let  $\varphi \in \text{End}(X)$   $\nearrow$  Frob. endom

•  $\varphi_* = \text{mult by } q^d$  on  $CH_d(X)$

$\varphi^* = \text{ " } q^m$  on  $CH^m(X)$

•  $P_m := \text{charpol}(\varphi^* \curvearrowright H^m(X))$

motivic arg.  $\Rightarrow P_m(\varphi^*) = 0$

on  $CH_{(s)}^i(X)$  if  $2i-s=m$

•  $q^i$  is a root of  $P_m$

Weil conj  $\Rightarrow m = 2i$   
 $s = 0$

3

Cor:  $X/\bar{\mathbb{F}}_p$  then:

$$CH_0(X) \cong \mathbb{Z} \oplus X(\bar{\mathbb{F}}_p)$$

Reason:  $CH_0 \cong \mathbb{Z} \oplus I$

$$\begin{array}{ccccccc} 0 & \rightarrow & I^{*2} & \rightarrow & I & \xrightarrow{S} & X(\bar{\mathbb{F}}_p) \rightarrow 0 \\ & & \parallel & & & & \\ & & \oplus & & CH_{0, (s)} & & \\ & & s \geq 2 & & & & \\ & & \parallel & & & & \\ & & 0 & & & & \end{array}$$

---

$Y/h = \bar{h}$  Sm. proj. var

$$CH_0(Y) \supset CH_0(Y)_{\text{hom}}$$

? What would it mean to say  
 $CH_0$  is "small"?

• Property A :

$\exists m$  s.t. every class in  $CH_0(Y)_{\text{hom}}$  can be repr'd as

$$P_1 + \dots + P_m - Q_1 - \dots - Q_m$$


---

•  $\gamma_m : Y^m \times Y^m \longrightarrow CH_0(Y)_{\text{hom}}$   
 $(P_1, \dots, P_m, Q_1, \dots, Q_m) \longmapsto P_1 + \dots + P_m - (Q_1 + \dots + Q_m)$

Fibres are countable unions of alg. subvars.

$$d_m = 2m \cdot \dim(Y) - \left( \begin{array}{l} \text{max. dim of a} \\ \text{subvar contained} \\ \text{in a fibre} \end{array} \right)$$

Property B

$m \mapsto d_m$  bounded

Property C :

$\exists$  non-singular curve  $C + j: C \rightarrow Y$

such that

$$j_* : CH_0(C)_{\text{hom}} \longrightarrow CH_0(Y)_{\text{hom}}$$

---

Prop. D Choose  $\gamma_0 \in Y(k)$

$$\text{alb} : CH_0(Y)_{\text{hom}} \longrightarrow \text{Alb}_Y(k)$$

$\text{alb}$  is an  $\cong$

Thm (A) - (D) are all equiv.

6

Theorem (Mumford 6g  
~ Roitman)

Suppose  $Y/\mathbb{C}$  is sm. proj. such  
that (A) - (D) hold.

Then  $H^0(Y, \Omega_Y^i) = 0 \quad \forall i \geq 2$

Cor  $X/\mathbb{C}$  Av,  $g \geq 2$

then (A) - (D) do not hold.

Thm (Bloch)  $k = \bar{k}$ ,  $\text{char} = 0$   
uncountable

$$X/k \quad \dim = g$$

$$I = CH_0(X)_{\text{hom}}$$

$$I^{*g} \neq 0 \quad (\text{recall: } I^{*(g+1)} = 0)$$

Eq: All  $CH_{0,(s)}$   $s = 0, \dots, g$   
are nonzero

False in  $\text{char} = p$  !

$X$  supersingular then all  $CH_{(s)}^i$   
 $s > 2$  vanish

"Complexity" of CH :

- Kimura / O'Sullivan :

fin. dim'l Chow motives

Know "motives of ab. type"

is fin. dim

- Voevodsky's Smash nilp. conj :

$\alpha \in CH^i(Y)$  if  $\alpha \equiv 0 \pmod{num}$  ( $\Leftarrow \alpha \sim_{hom} 0$ )

then  $\exists N :$

$\alpha^{\otimes N} : \underbrace{\alpha \times \alpha \times \dots \times \alpha}_N \in CH(Y^N)$

$\parallel$

0

Known if  $\alpha \sim_{alg} 0$

Known for 1-cycles on AV